



DiscoverSys  
Whatever it takes...

Published by DiscoverSys

# Anti tuberculosis activity of forest Kedondong (*Spondias pinnata*) stem bark extract against Multiple Drug Resistance (MDR) strain of *Mycobacterium tuberculosis*



CrossMark

I.B.N. Putra Dwija,<sup>1</sup> Mita Anggraeni,<sup>2</sup> Ni Putu Ariantari<sup>2\*</sup>

## ABSTRACT

**Background:** Forest Kedondong (*Spondias pinnata*) traditionally known as “*loloh cemcem*” and commonly used as a chronic cough remedy. Previous research showed that methanol extract of Forest Kedondong leaves active against MDR strain of *Mycobacterium tuberculosis*. The aim of this study were to determine the phytochemical constituent and anti tuberculosis activity of stem bark extract of this plant against MDR strain of *M. tuberculosis*.

**Method:** Coarsely powder of Forest Kedondong stem bark was extracted successively with n-hexane, chloroform and 80% ethanol. Anti tuberculosis assay of chloroform and ethanol extract was conducted using proportion method with Lowenstein-Jensen medium within 3 different concentration of extract of 1, 10, and

100 mg/mL. Activity was evaluated by inhibition of extract against *M. tuberculosis* growth, which was calculated by mean reduction in number of colonies on extract containing medium compared to control.

**Results and Discussion:** Phytochemical test showed that chloroform extract contains terpenoid and flavonoids. Ethanol extract contains terpenoid, polyphenols and flavonoids. These extracts were active against MDR strain of *M. tuberculosis* with 100% inhibition at concentration of 100 mg/mL. Chloroform extract has higher inhibition against *M. tuberculosis* growth than Ethanol extract.

**Conclusions:** These extracts were potentially developed to an anti tuberculosis constituent from natural product.

**Keywords:** Kedondong stem bark, chloroform extract, Ethanol extract, *Mycobacterium tuberculosis*, anti tuberculosis.

**Cite This Article:** Dwija, I., Anggraeni, M., Ariantari, N. 2016. Anti tuberculosis activity of forest Kedondong (*Spondias pinnata*) stem bark extract against Multiple Drug Resistance (MDR) strain of *Mycobacterium tuberculosis*. *Bali Medical Journal* 5(1): 23-26. DOI: [10.15562/bmj.v5i1.190](https://doi.org/10.15562/bmj.v5i1.190)

<sup>1</sup>Department of Microbiology, Faculty of Medicine, Udayana University, Bali-Indonesia

<sup>2</sup>Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Udayana University

## INTRODUCTION

Tuberculosis is infectious disease caused by *Mycobacterium tuberculosis*, which mostly attack lungs. According to WHO report (2013), Indonesia was become the 3<sup>rd</sup> country with high prevalence of tuberculosis in Asia after India and China. Pandemic of HIV/AIDS significantly increase tuberculosis cases all over the world. Mortality of tuberculosis more frequent in poor and developing countries. Another problem arise in combatting tuberculosis is the presence of resistant strain of *M. tuberculosis*, including mono resistant, multi-drug resistant and extended drug resistant strains of *M. tuberculosis*. Treatment of multidrug resistant tuberculosis need higher cost and possibly cause more side effects and drug toxicity. Therefore, many research has been conducting to develop new anti tuberculosis agent.<sup>1</sup>

Medicinal plants have been proven to be an important resource of many therapeutic agents. WHO (2003) also recommended the usage of ethno-medicinal plants in health prevention, promotion and curative of disease. *Spondias pinnata* which is locally known as Forest kedondong, was

used as traditional drink in Bali known as “*loloh cemcem*”.<sup>2</sup> Leaves of this plant also traditionally used for chronic cough remedy. Forest kedondong belongs to Anacardiaceae family. Methanol extract of Forest kedondong leaves showed potent inhibition against multidrug resistant (MDR) strain of *M. tuberculosis in vitro*.<sup>3</sup> N-hexane extract also revealed inhibition against MDR strain of *M. tuberculosis*.<sup>4,5</sup> Extract of *S. pinnata* fruit revealed cytotoxic effect and antibacterial activity against *Pseudomonas aeruginosa* and *Staphylococcus epidermidis*.<sup>6</sup> Extract of *S. pinnata* stem bark also reported for its anthelmintic activity and antioxidant.<sup>7,8</sup> In present research we evaluate anti tuberculosis activity of *S. pinnata* stem bark extract against MDR strain of *M. tuberculosis*.

## MATERIAL AND METHODS

### Plant material

Forest kedondong stem bark was collected from Badung, Bali-Indonesia. Identification of plant specimen was conducted at Kebun Raya Eka Karya, Bedugul, Tabanan-Indonesia.

\*Correspondence to: Ida Bagus Putra Dwija, Department of Microbiology, Faculty of Medicine, Udayana University, Bali-Indonesia. [ari\\_dedhika@yahoo.com](mailto:ari_dedhika@yahoo.com)

### Bacteria *M. tuberculosis* MDR

Bacteria was obtained from Clinical Microbiology Department, Sanglah General Hospital, Denpasar, Bali-Indonesia.

### Extraction

Coarsely dried powder of Forest kedondong stembark (50 g) was extracted successively with n-hexane, chloroform and 80% of ethanol to obtain chloroform extract (2.50 g) and ethanoic extract (5.05 g).

### Phytochemical test

Phytochemical test was conducted according to methods previously described by Evans et al. (2009), Jones and Kinghom (2006) and Depkes RI (1989) to detect the presence of alkaloids (Dragendorff reagent), saponins (ability to produce stable foam), polyphenols (ferric chloride reagent), steroid, terpenoids and glycosides (Lieberman-Bouchard reagent), flavonoids (with the use of boric acid and oxalic acid).<sup>9-11</sup>

### Activity Assay

Anti tuberculosis activity assay was conducted using proportion method, according to Gupta et al. (2010).<sup>12</sup> Extract was added into Lowenstein-Jensen medium to give serial concentration of extract of 1, 10, and 100 mg/mL. Control group only received 1% dimethyl sulfoxide. Bacteria then inoculated on this medium, incubated in 5% CO<sub>2</sub> incubator, 37°C for 6 weeks. Observation of colonies growth

was done starting from 3<sup>rd</sup> until 6<sup>th</sup> week. Inhibition of extract against *M. tuberculosis* was calculated by comparing colonies growth in treatment group which received series concentration of extract to control.

## RESULTS AND DISCUSSION

In extraction process, ethanol solvent gave the higher yield than chloroform. N-hexane is used to remove non polar soluble components like resins and other miscellaneous compounds. Phytochemical test showed that chloroform extract contains terpenoids and flavonoids and ethanol extract contain terpenoids, flavonoids and polyphenols. The forming of brown color ring indicated presence of terpenoids and flavonoids were represented by appearance of intensive yellow fluorescence under UV light 366 nm. The color changes of ethanoic extract into dark green color indicated the presence of polyphenols.


The result of anti tuberculosis activity assay of extracts was shown in table 1.

Observations were made from 3<sup>rd</sup> to 6<sup>th</sup> week because this period is growth phase of *M. tuberculosis* on L-J medium.<sup>13</sup> Increasing concentration of chloroform and Ethanol extract on medium resulting in higher inhibition to *M. tuberculosis* growth. Comparison of inhibition of each extracts was shown in figure 1.

At the extract concentration of 1 and 10 mg/mL, chloroform extract gave higher inhibition than

**Table 1** Inhibition of extract of Forest Kedondong stembark against MDR strain of *M. tuberculosis*

Observation	Inhibition against <i>M. tuberculosis</i> growth (%)						
	Extract concentration (mg/mL)						
		Chloroform extract			Ethanoic extract		
		1	10	100	1	10	100
3 <sup>rd</sup> week	1	25.00	75.00	100	12.50	68.75	100
	2	35.29	84.31	100	29.41	80.39	100
	3	51.96	82.35	100	49.02	80.39	100
4 <sup>th</sup> week	4	57.24	83.45	100	55.86	82.07	100
	5	51.15	85.06	100	50.00	83.91	100
	6	52.11	84.39	100	50.24	81.46	100
5 <sup>th</sup> week	7	49.79	84.55	100	48.41	82.83	100
	8	44.17	83.33	100	42.50	82.50	100
	9	41.77	82.33	100	40.56	81.53	100
6 <sup>th</sup> week	10	39.92	80.24	100	38.34	78.66	100
	11	39.39	79.55	100	37.88	77.27	100
	12	39.39	79.55	100	37.88	77.27	100



**Figure 1** Comparison of inhibition of chloroform (EK) and Ethanol extract (EE) of Forest kedondong stem bark against *M. tuberculosis* growth at extract concentration of 1 mg/mL (A), 10 mg/mL (B) and 100 mg/mL (C)

Ethanol extract (Figure 1 (A) and (B)). Treatment with extract at concentration of 100 mg/mL, which is shown in Figure 4.1 (C), chloroform and Ethanol extract of Forest Kedondong stem bark able to inhibit growth of *M. tuberculosis* during the observation period by inhibition of 100%. According to Gupta et al. (2010), the extract is stated to have anti tuberculosis activity when inhibition to bacteria growth is  $\geq 90\%$ . Based on these criteria, chloroform and Ethanol extract of Forest kedondong stem bark had anti tuberculosis activity against MDR strain of *M. tuberculosis* at concentration of 100 mg/mL.<sup>12</sup>

Various studies on anti tuberculosis activity of flavonoid and terpenoids also been reported. Flavonollaburnetin isolated from *Ficus cordata* and *Ficus chlamydocarpa* also showed anti tuberculosis activity with MIC value of 4.88  $\mu\text{g/mL}$  (Kuethe, 2008). Isoflavonoids isolated from the roots of *Sesbaniagrandiflora* showed anti tuberculosis activity against *M. tuberculosis H37Rv* with MIC of 50  $\mu\text{g/mL}$  (Hasan, et al., 2012). According to Brown et al. (2007), many flavonoids can inhibit fatty acid synthase type II which involving in mikolat acid biosynthesis, component of cell wall of *M. tuberculosis*. Olugbuyiro et al. (2009) reported triterpenoid compounds derived from methanol extract of *Spondias mombin* L. bark (Anacardiaceae family) has anti tuberculosis activity against *M. tuberculosis H37Rv* (sensitive strain) at extract concentration of 64  $\mu\text{g/mL}$ .<sup>14-17</sup>

## CONCLUSION

Both chloroform and Ethanol extract of Forest kedondong stem bark were active as anti

tuberculosis against MDR strain of *M. tuberculosis* with 100% inhibition at concentration of 100 mg/mL. Terpenoids and flavonoids might be contributed to the activity of these extracts. However, further research is suggested to identify anti tuberculosis constituents from these extracts.

## REFERENCES

- World Health Organization. (2013). *Global Tuberculosis Report 2013*. France: World Health Organization. P. 19, 61.
- World Health Organization. (2003). *Traditional medicine*. (serial online), (cited 2014 Okt, 5). Available from: <http://www.who.int/mediacentre/factsheets/fs134/en/>.
- Dwija, I.B.N.P., Juniarta, I.K., Yowani, S.C., and Ariantari, N.P. (2013). Aktivitas Anti tuberculosis Ekstrak Metanol Daun Kedondong Hutan (*Spondiaspinnata* (L.F) Kurz.). *Jurnal Kimia*. Vol. 7 (1): 25-30
- Savitri, L. P. V. A., Ariantari, N. P., and Dwija, I. B. N. P. (2013). Potensi Anti tuberculosis Ekstrakn-heksana Daun Forest kedondong (*Spondiaspinnata* (L.f) Kurz.). *Jurnal Farmasi Udayana*. Vol. 2 (3): 105-109
- Ramayati, N. P. A., Ariantari, N. P., and Dwija, I. B. N. P. (2013). Aktivitas Anti tuberculosis Kombinasi Ekstrak n-heksana Daun Forest kedondong dengan Rifampisin Terhadap Isolat *Mycobacterium Tuberculosis* Strain MDR. *Jurnal Farmasi Udayana*. Vol. 2 (3): 74-78
- Muhammad, A., Rahman, M. S., Kabir, A. H., Kabir, S., and Hossain, M. K. (2011). Antibacterial and cytotoxic activities of *Spondiaspinnata* (Linn.f) Kurz fruit extract. *Indian Journal of Natural Products and Resources*, Vol. 2 (2): 265-267.
- Panda, B.K., Patro, V.J., Mishra, U.S. and Panigrahi, B.K. (2011). Comparative Study of Anthelmintic Activity between Acetone and Ethanol Stem Bark Extracts of *Spondiaspinnata* (Linn.F) Kurz. *International Journal of Research in Ayurveda & Pharmacy*. Vol. 2(4):1383-1385
- Hazra, B., Biswas, S., Mandal, N., (2008). Antioxidant and Free Radical Scavenging activity of *Spondiaspinnata*. *BMC Complementary and Alternative Medicine*. Vol. 8:63
- Evans, C. W. (2009). *Pharmacognosy Trease and Evans 16<sup>th</sup> Edition*. China: Saunders Elsevier. P. 263-356.
- Jones, W. P. and Kinghorn, A. D. (2006). Extraction of Plant Secondary Metabolites. In: Sarker, S. D., Latif, Z. and Gray, A. I., editors. *Natural Products Isolation*, Second Edition. New Jersey: Humana Press. P. 341-342

11. Depkes RI. (1989). *Materia Medika Indonesia V*. Jakarta: DepartemenKesehatanRepublik Indonesia: Hal: 549-553
12. Gupta, R., Thakur, B., Singh, P., Singh, H.B., Sharma, V.D., Katoch, V.M., & Chauhan, S.V.S. (2010). Anti-tuberculosis Activity of Selected Medicinal Plants Against Multi-Drug Resistant *Mycobacterium Tuberculosis* Isolates. *International Journal Medicine Research*, Vol. 131: 809-813.
13. Todar, K. (2008). *Textbook of Bacteriology* (serial online), (cited: 2013 Feb, 20). Available from: <http://www.textbook-bacteriology.net/tuberculosis.html>
14. Brown, A.K., Papaemmanouil, A., Bhowruth, V., Bhatt, A., Dover, L. G. and Besra, G. S. (2007). Flavonoid inhibitors as novel antimycobacterial agents targeting Rv0636, a putative dehydratase enzyme involved in *Mycobacterium tuberculosis* fatty acid synthase II. *Microbiology*, Vol. 153: 3314-3322.
15. Hasan, N., H. Osman, M. Suriyanti., W. K. Chong, K. Awang and A. S. M. Zahariluddin. (2012). The Chemical Components of *Sesbaniagrandiflora*Root and Their Anti Tuberculosis Activity. *Journal of Pharmaceuticals*. Vol. (5): 882-889
16. Kuete, V., Ngameni, B., Fotso-Simo, C.C., Kengap, T.R., Tchaleu, N.B., Meyer J.J.M., Lall N., danKuiate, J.R. (2008). Antimicrobial activity of the crude extracts and compounds from *Ficuschlamydocarpa* and *Ficuscordata* (Moraceae). *J. Ethnopharmacol*, Vol. 120: 17-2.
17. Olugbuyiro, J. A. O., Mody, J. O., and Hamann, M. T. (2009). AntiMtb Activity of Triterpenoid-Rich Fractions From *Spondiasmombin* L. *African Journal of Biotechnology*, Vol. 8 (9): 1807-1809.



This work is licensed under a Creative Commons Attribution